



+ + + +

# Diagnosing intestinal parasite infections

Fecal Dx<sup>®</sup> antigen testing clinical reference guide

**IDEXX**

# Screen dogs and cats at least twice a year

The Companion Animal Parasite Council (CAPC) guidelines recommend including fecal antigen testing to ensure the widest breadth of detection of intestinal parasites.<sup>1-3</sup> Fecal antigen testing can identify infections that can be missed by using other methods.<sup>4</sup>

## Hidden dangers and recommendations pet parents should know.

CAPC recommends fecal screening at least twice a year for adult dogs and cats but at least four times during the first year of a pet's life. A pet's health and lifestyle may warrant more frequent testing. Simply being outside can put pets at risk. Some parasites can burrow into a pet's skin or feet. Dogs and cats can also swallow parasites while grooming, nursing, or eating contaminated soil or feces. Clients should also know that some infections can spread from pets to people.<sup>5</sup>

## Recommend year-round, broad-spectrum parasite control.

A parasite-control program that is effective against intestinal parasites, heartworms, ticks, and fleas provides maximum value for your clients and the best protection for your patients. Make sure your clients understand that they need to keep their pets protected all year long.<sup>5</sup>

## Diagnose and treat infections earlier with Fecal Dx<sup>®</sup> antigen testing.

CAPC recommends including fecal antigen testing to diagnose infections, treat patients earlier, and reinforce the proper use of parasite-control products.<sup>1-3</sup> Fecal antigen testing identifies prepatent and single-sex infections, providing critical insights for patient management.<sup>1-3</sup>




# Fecal screening for healthy adult pets

Diagnose hookworm, roundworm, whipworm, flea tapeworm, and *Cystoisospora* infections using Fecal Dx<sup>®</sup> antigen testing. Because the testing detects antigens, positive results confirm the presence of worms or protozoa in the intestinal tract. This allows you to diagnose infection when worms are not shedding eggs or are caused by worms of a single sex.<sup>1-3</sup>

## What to do with your Fecal Dx antigen testing results.

The following algorithm can help guide your next steps when considering your patient's Fecal Dx antigen testing results.



# Did you know?

Treatment should be considered for patients that test positive by either antigen testing or egg/proglottid and oocyst detection.

Reasons for specimens that are antigen positive and egg/proglottid/oocyst negative may include the following:

- + Absence of eggs, proglottids, and oocysts during the prepatent period
- + Single-sex infections (worms)
- + Intermittent egg/proglottid/oocyst shedding

Reasons eggs, proglottids, and oocysts may be identified in specimens that are antigen negative may include the following:


- + Ingestion of infected feces (coprophagy)
- + Antigen quantity is below the level of detection



# Hookworms: from infection to presentation<sup>1</sup>

## *Ancylostoma caninum* life cycle

Prepatent period for adult dogs: 14–21 days



## Did you know?

- + Because hookworms have short prepatent periods and the potential for arrested larvae, even pets receiving monthly deworming may have adult worms in their intestinal tract between monthly doses.<sup>1</sup>
- + Resistance to dewormers is now documented for *A. caninum*.<sup>6</sup> If hookworm positive 10–14 days following deworming, a fecal egg count reduction test should be performed.<sup>7</sup>
- + Puppies as young as 10–12 days of age may start shedding eggs if they've been infected through nursing.<sup>1</sup>
- + Due to the zoonotic risk and reinfection potential in this parasite, it is important to detect infections before they start shedding eggs into the environment.<sup>1</sup>


## Clinical presentation

Pale mucous membranes and anemia; ill thrift, failure to gain weight; poor hair coat, dehydration; dark, tarry diarrhea; respiratory disease; foot lesions (dermatitis with erythema, pruritus, and papules).

# Roundworms: from infection to presentation<sup>2</sup>

## *Toxocara canis* life cycle

Prepatent period for adult dogs: 21–35 days



### Did you know?

- + One female roundworm can produce 85,000 eggs per day, and these hard-shelled eggs can survive in the environment for years.<sup>2</sup> With Fecal Dx<sup>®</sup> antigen testing, you can detect infections before roundworms start laying eggs.
- + Due to the zoonotic risk and reinfection potential in this parasite, it is important to detect infections before they start shedding eggs into the environment.<sup>2</sup>
- + In puppies under 6 months of age, studies have shown more than 30% are infected and shedding *T. canis* eggs.<sup>2</sup>



### Clinical presentation

Diarrhea, vomiting, pot-bellied appearance, coughing. Dogs may cough up or vomit worms. Infections caused by *T. canis* are more common and most severe in dogs less than one year of age.

# Whipworms: from infection to presentation<sup>3</sup>

## *Trichuris vulpis* life cycle

Prepatent period for adult dogs: 74–90 days



## Did you know?

- + A female whipworm can produce as many as 2,000 eggs per day, and these infective whipworm eggs can survive in the environment for several years.<sup>3</sup> With Fecal Dx<sup>®</sup> antigen testing, you can detect infections before whipworms start laying eggs.
- + Due to their extended prepatent period, it's unlikely to find eggs being shed in very young puppies,<sup>3</sup> but Fecal Dx antigen testing can identify these positive patients during the prepatent period for earlier diagnosis and treatment.

## Clinical presentation



Many infections are subclinical. When present, clinical signs include diarrhea streaked with mucus and fresh blood, weight loss, dehydration, anemia. Extreme cases can result in death.

# Flea tapeworm: from infection to presentation<sup>8</sup>

## *Dipylidium caninum* life cycle

Prepatent period for adult dogs: 14–35 days<sup>8,10</sup>

This tapeworm can infect dogs and cats and is spread through ingestion of infected fleas.



## Did you know?

- + *D. caninum* is called the “flea tapeworm” because the flea is its intermediate host.<sup>8</sup>
- + Dogs and cats become infected by eating/ingesting an infected flea.<sup>8</sup>
- + Each segment (proglottid) of a flea tapeworm may contain up to 25–30 eggs.<sup>8</sup>
- + Reinfection with *D. caninum* is likely if flea infestations are not controlled.<sup>8</sup>
- + Dogs and cats may be infected with more than one species of tapeworm. The flea tapeworm is most commonly diagnosed.<sup>9</sup>

## Clinical presentation

Infection may not always be apparent as many are subclinical. While flea tapeworms rarely cause disease, the passage of the proglottids may cause perianal irritation.




# Cystoisospora spp. from infection to presentation<sup>11</sup>

## Cystoisospora spp. life cycle

Prepatent period 4–13 days<sup>11</sup>

This protozoa can infect dogs and cats and is spread through ingestion of sporulated oocysts in the environment or predation of an infected paratenic host.

Dog ingests sporulated oocysts  
or infected tissue of paratenic host



## Did you know?<sup>1</sup>


- + Oocysts may survive up to a year in moist, protected conditions if freezing or excessively high temperatures are avoided.<sup>11</sup>
- + Oocysts of a pseudoparasite, *Eimeria* spp. are sometimes found in dog and cat feces.<sup>11</sup> Their morphology is very similar to *Cystoisospora*.
- + *Eimeria* is a coccidian parasite of rabbits, birds, rodents, and reptiles/amphibians but does not cause infection or disease in dogs and cats.<sup>11</sup>

## Clinical presentation

Healthy adult dogs and cats may be infected and show no clinical signs. Young, stressed, and/or immunocompromised patients may have diarrhea, vomiting, dehydration, depression, and anorexia.


# Fecal screening for sick dogs and cats


When assessing sick pets with diarrhea, it is important to differentiate between simple and complicated cases. For simple diarrhea of short duration, testing for *Giardia* antigen by immunoassay or the SNAP® *Giardia* Test, as well as Fecal Dx® antigen testing and fecal ova and parasites by centrifugation (fecal O&P) is recommended. In cases of complicated diarrhea, a more comprehensive diarrhea RealPCR™ panel is recommended to rule out infectious diseases.



## Did you know?

For complicated sick patients, our RealPCR™ testing expands the scope of detection to other important gastrointestinal pathogens, including viruses, bacteria, enterotoxins, and difficult-to-detect protozoa (e.g., *Cryptosporidium*).





# How to minimize the threat of parasite infection

In addition to regular veterinary examinations and diagnostic screening, encourage your clients to follow CAPC guidelines.<sup>5</sup>

## Food and water


- + Pets should be fed commercial or cooked food.
- + Raw diets are not recommended.
- + Make sure pets have access to fresh water.

## Outdoor activity

- + Limit access to wildlife and contaminated feces.
- + Keep dogs on a leash or behind a fence.
- + Keep cats indoors.
- + Cover sandboxes after use.

## Hygiene

- + Do not handle animal feces or urine with bare hands.
- + Wash hands immediately after incidental contact with feces.
- + Promptly pick up and dispose of feces from public areas.
- + Promptly remove feces from the yard.





## References

1. Companion Animal Parasite Council. CAPC guidelines: hookworms. Updated March 29, 2023. Accessed February 19, 2024. [www.capcvet.org/guidelines/hookworms](http://www.capcvet.org/guidelines/hookworms)
2. Companion Animal Parasite Council. CAPC guidelines: ascarid. Updated September 12, 2022. Accessed February 19, 2024. [www.capcvet.org/guidelines/ascaris](http://www.capcvet.org/guidelines/ascaris)
3. Companion Animal Parasite Council. CAPC guidelines: *Trichuris vulpis*. Updated September 12, 2022. Accessed February 19, 2024. [www.capcvet.org/guidelines/trichuris-vulpis](http://www.capcvet.org/guidelines/trichuris-vulpis)
4. Adolph C, Barnett S, Beall M, et al. Diagnostic strategies to reveal covert infections with intestinal helminths in dogs. *Vet Parasitol.* 2017;247:108–112. doi:10.1016/j.vetpar.2017.10.002
5. General guidelines for dogs and cats. Companion Animal Parasite Council website. Accessed July 13, 2022. [www.capcvet.org/guidelines/general-guidelines](http://www.capcvet.org/guidelines/general-guidelines)
6. Jimenez Castro PD, Howell SB, Schaefer JJ, Avramenko RW, Gilleard JS, Kaplan RM. Multiple drug resistance in the canine hookworm *Ancylostoma caninum*: an emerging threat? *Parasit Vectors.* 2019;12(1):576. doi:10.1186/s13071-019-3828-6
7. Jimenez Castro PD, Kaplan RM. Persistent hookworm infections in dogs. *Clin Brief.* August 2020;59. Accessed February 19, 2024. [www.cliniciansbrief.com/article/persistent-hookworm-infections-dogs](http://www.cliniciansbrief.com/article/persistent-hookworm-infections-dogs)
8. CAPC guidelines: *Dipylidium caninum*. Companion Animal Parasite Council website. Accessed February 19, 2024. [www.capcvet.org/guidelines/dipylidium-caninum](http://www.capcvet.org/guidelines/dipylidium-caninum)
9. Hall EJ, Day MJ. Diseases of the small intestine. In: Ettinger SJ, Feldman EC, Côté E, eds. *Textbook of Veterinary Internal Medicine: Diseases of the Dog and Cat.* 8th ed. Elsevier; 2017:1543.
10. Beugnet F, Labuschagne M, Vos C, Crafford D, Fourie J. Analysis of *Dipylidium caninum* tapeworms from dogs and cats, or their respective fleas—Part 2. Distinct canine and feline host association with two different *Dipylidium caninum* genotypes. Analyse des ténias *Dipylidium caninum* des chiens et des chats, ou de leurs puces respectives - Partie 2. Association distincte des hôtes canins et félins avec deux génotypes différents de *Dipylidium caninum*. *Parasite.* 2018;25:31. doi:10.1051/parasite/2018029
11. Companion Animal Parasite Council. CAPC Guidelines: Coccidia. Updated October 1, 2016. Accessed February 19, 2024. [www.capcvet.org/guidelines/coccidia](http://www.capcvet.org/guidelines/coccidia)
12. Lappin MR. Isosporiasis. In: Sykes JE, ed. *Canine and Feline Infectious Diseases.* WB Saunders; 2014:793–796. doi:10.1016/B978-1-4377-0795-3.00082-X

© 2024 IDEXX Laboratories, Inc. All rights reserved. • 09-81850-04  
All ®/TM marks are owned by IDEXX Laboratories, Inc. or its affiliates in the United States and/or other countries.  
The IDEXX Privacy Policy is available at [idexx.com](http://idexx.com).

PCR testing is a service performed pursuant to an agreement with Roche Molecular Systems, Inc.