



# Oh no otitis!


## When to reach for antimicrobials, which one to choose and more

Christina Gentry, DVM DACVD  
Assistant Clinical Professor of Veterinary Dermatology  
Texas A&M University, College Station, Texas, USA



# Financial Disclosure

I have a commercial relationship with IDEXX company, but do not believe that it will influence my presentation.



# Learning Objectives

- **Define key terminology** related to ear canal changes in otitis, including erythema, stenosis, hyperplasia, ceruminous discharge, and purulent discharge
- **Analyze case examples of progressive otitis externa** to inform therapeutic decision-making.
- **Select appropriate antimicrobial and anti-inflammatory therapies** based on disease progression and presentation.
- **Recognize anatomic variations in brachycephalic breeds** and adapt treatment strategies to optimize outcomes in these patients.

# The ear examination

Palpate the ear canals on both sides



- First/Second time offenders should have soft and pliable canals
- Chronic offenders may have firm to hard (calcification) ear canals
- Chronic offenders can have nodular hyperplasia (large bumps) in the ear canal that make the canal thick and non pliable

Evaluate the pinna and entrance of the canal



- Are the pinna affected?
- Look for nodular hyperplasia at the entrance of the ear canal

# Otoscopic examination

- Pull the pinna towards you and down slightly.
- Your head should be at the same or just above the height of the patient ear




# The normal ear canal

The normal ear canal is temperature, pH, and humidity regulated.

Normal ear canals rely on the lipid content of cerumen, host defense(antimicrobial) peptides, and regulated desquamation to maintain homeostasis.

The ear canal is light pink, smooth, and has minimal debris.




# Impression cytology of the normal ear

Unstained samples will appear nearly clear and slightly greasy

After staining it is still nearly clear since the lipid does not take up much stain

Few anucleated keratinocytes(squamous cells) will pick up purple stain

\*\*\*A few yeast or cocci adhered to squamous cells is within the realm of normal



100x objective, few keratinocytes


# Erythematous ear canals



Erythema, stenosis



Erythema,  
stenosis, nodular  
hyperplasia




Erythema, stenosis,  
epithelial  
hyperplasia

# Ear canal wall changes- epithelial hyperplasia



Minimally erythematous smooth walls




Mild erythema and mild epithelial hyperplasia with stenosis




Mild erythema and moderate epithelial hyperplasia with stenosis

# Ear Canal wall changes-nodular hyperplasia




Irregular nodular hyperplasia and stenosis



Large nodules obstructing the canal

# Description of debris



Mild to moderate ceruminous debris, obstructs TM



Heavy ceruminous debris with ceruminolith formation



Moderate purulent debris



Slimy purulent debris with biofilm formation

# Tympanic Membrane of the Dog and Cat

The tympanic membrane (tympanum, TM) is external border of the middle ear cavity.

**The slight concavity of the pars tensa is due to the insertion of the manubrium of malleus attachment** under the epithelium applying slight pressure to the tympanum

(This part of the TM is easiest to visualize in the normal dog)

The pars flacida is confined to the upper quadrant  
(This part of the TM is harder to visualize in the normal dog)



Feline Tympanic Membrane with manubrium of malleus visible

# Tympanic membrane in the dog



Clear TM with hair tuft attached to adjacent canal wall




Slightly opaque but fully visible TM



Opaque TM is partially covered by ceruminous debris

# Evaluation of ear drums- Tympanic membrane rupture



Intact canine tympanic membrane



Ruptured canine tympanic membrane

# *Malassezia* Yeast

## *Malassezia pachydermatis*

Opportunistic pathogen that is part of the normal flora

Bowling pin, snowman, peanut shaped when budding


Stain deep blue/purple with Diff Quick®

### ***Treatment:***

Terbinafine

Azoles (Posaconazole, thiobendazole, miconazole, clotrimazole)

*Corticosteroid Ototopical*



# Cocci bacteria

*Staphylococcus pseudintermedius*

Opportunistic pathogen that is part of the normal flora

Deeply blue/purple perfect circles when stained with Diff Quick®


## ***Treatment:***

Aminoglycoside (gentamicin, neomycin)

Florfenicol

+/-Fluroquinolone (enrofloxacin)

*Corticosteroid Ototopical*



# Rod Shaped bacteria

## *Pseudomonas aeruginosa*

- Rod shaped bacteria are never 'normal' on otic cytology
- Deeply blue/purple rods
- Pseudomonas* requires inflammation + predisposing factor (increased humidity, increased pH) to overgrow
- Rod shaped otitis is more common in ears that have chronic changes from previous inflammatory and infectious (cocci, yeast) insults.


### **Treatment:**

Polymixin B

Fluoroquinolone (enrofloxacin)

Silver Sulfadiazine

Aminoglycoside (Amikacin)



# Treatment of otitis externa- TM intact

## Yeast Otitis

EasOtic<sup>®</sup>  
Mometamax<sup>®</sup>

Leave ins  
Claro<sup>®</sup>, Osurnia<sup>®</sup>,  
Mometamax Single<sup>®</sup>,  
DuOtic<sup>®</sup>

Stand alone topical  
corticosteroid

## Cocci Otitis

EasOtic<sup>®</sup>  
Mometamax<sup>®</sup>

Leave ins  
Claro<sup>®</sup>, Osurnia<sup>®</sup>  
Mometamax Single<sup>®</sup>

Stand alone topical  
corticosteroid

## Rod shaped otitis

Surolan<sup>®</sup>  
Baytril Otic<sup>®</sup>

Mometamax Single<sup>®</sup>  
(*Pseudomonas*)

# Treatment of otitis externa- unknown TM status/ TM ruptured

## Yeast Otitis

Miconazole  
/Dexamethasone

## Cocci Otitis

Enrofloxacin/  
Miconazole/  
Dexamethasone  
in Triz-ETDA  
base

## Rod shaped otitis

Enrofloxacin/  
Miconazole/  
Dexamethasone  
in Triz-ETDA  
base

# Ear Flushing and Otitis

## ***Ceruminous Otitis Externa***

### **Products that are oil based or feel 'oily'**

Squalene- Cerumene®

pH Notix®

Douxo® Micellar Solution

### **Products that are drying**

Salicylic acid – EpiOtic Advanced®

Malacetic Ultra®

Cort/Astrin® Solution

## ***Purulent***

### **Antibacterial agents**

Isopropyl alcohol

Chlorhexidine

Hypochlorous acid

### **Adjunctive Antimicrobial**

Triz- EDTA

Monosaccharides,  
polysacaccharides

# Ready for case examples?

.....probably want to put your breakfast away.....

# Case Presentation -2 year old GSD Mix


## First Time Offender

HX: Went swimming in the pond twice last weekend and is now shaking head.

No prior ear infection, mild pedal pruritus in the spring and summer, monthly isoxazoline flea/tick control. Presumed healthy animal.

PE: No skin erythema, no ectoparasites, BCS 5/9

Both ear canals similarly affected. TM are intact



Minimal erythema, moderate dark brown waxy debris

# Case Presentation- 2 year old GSD Mix

## First Time Offender

### *Treatment:*


Corticosteroid ototopical and cleanser

OR leave in terbinafine + antibacterial ototopical

OR azole ototopical + antibacterial ototopical

### **Do I clean this dog's ears before sending it home?**

Honestly, no –unless it's recommended prior to placing a leave in product



Moderate cocci and fewer yeast

# When to flush ear canals?

## **Prior to treatment**

*In ceruminous otitis with heavy debris*

If recommended by the leave in product manufacturer

## **During treatment**

*Ceruminous otitis – 1-2x weekly*

Purulent otitis- 3-4 times weekly  
(may be daily if your antibiotic is compounded into a flush solution)

# First Time Offender


Recheck 2 weeks later

HX: Applied medications as directed, head shaking has resolved

PE: No skin erythema, no ectoparasites, BCS 5/9

Ear canals have minimal erythema, but otherwise normal on otoscopic examination

***Treatment: monitor for new clinical signs, cerumenolytic and antimicrobial cleanser after swimming***



Few anucleate keratinocytes

# When to flush ear canals- after treatment

If there is underlying allergy leading to ceruminous build up

- Consider flushing canals 1-2 times per week with a flush that also contains a topical steroid

If there is underlying sebaceous adenitis leading to dry wax build up

- Consider flushing canals 1-2 times per week with a flush that hydrates the ear canals

# Case Presentation- 2 year old Bull Terrier

## Repeat Offender

HX: Went to the ranch for the weekend and is head shaking and digging at the ear. Four prior ear infections, moderate spring-fall pruritus of paws and trunk, monthly isoxazoline flea/tick control.

PE: Salivary staining on paws, excoriations on the axilla with occasional papules on axilla and inguinal region , no ectoparasites, BCS 5/9

Both ear canals similarly affected. TM are intact



Erythema, epithelial hyperplasia, ceruminous debris, ceruminolith

# Case Presentation – 2 year old Bull Terrier

## Repeat Offender

### ***Topical Plan***


Corticosteroid ototopical and cerumenolytic cleanser

OR azole antifungal ototopical + topical antibiotic and ceruminolytic cleanser

OR Terbinafine ototopical + topical antibiotic and ceruminolytic cleanser

### ***Systemic Plan***

Oral corticosteroid 0.5-1mg/kg per day until resolution of inflammatory changes, then taper



Moderate cocci and fewer yeast

# Why oral corticosteroids?

- Oral steroids are best to reduce ear canal inflammation and ear canal discomfort
  - Improves medication compliance
  - Reduces risk of immediate relapse
- Oral steroids can prevent and sometimes reverse chronic changes to the canal
  - Decrease hyperplastic change that increase risk of infection
  - Decrease risk of progression to ulcerated and purulent otitis

# How is the oral steroid dosed?

Moderate to high anti-inflammatory is continued until the patient is showing significant improvement.

0.5-1mg/kg q 24 hours for 7-14 days

0.5-1mg/kg q 48 hours for 7-14 days

0.5mg/kg q 72 hours for 7-14 days


# Case Presentation- 2 year old Bull Terrier

Recheck 2 weeks later

HX: Owner administered medications as directed, head shaking is improved but not resolved

PE: Reduced salivary staining on paws, papules and excoriations resolved.

Ear examination shows minimal debris with erythematous slightly irregular ear canals and focal stenosis



Erythema, epithelial and glandular hyperplasia, mild stenosis

# Case Presentation - 2 year Bull Terrier

## Maintenance Plan

### ***Topical Plan***

Otological steroid daily until inflammation resolves +/- a few times per week longer term

Ceruminolytic and antimicrobial flush 1-2 times per week and after swimming


### ***Systemic Plan***

**Slowly taper oral steroids**

Modified cyclosporine

Oclacitinib or Ilunocitinib

Allergy specific immunotherapy (+ CsA or oclacitinib)



Variably stained keratinocytes

# Which otic topical corticosteroids is right for this patient?

- Lower potency steroids for infrequent or mild inflammation
  - Hydrocortisone
- Higher potency steroids for frequent otitis or moderate inflammation
  - Mometasone (less systemic absorption)
  - Injectable dexamethasone (moderate systemic absorption)
  - Fluocinolone acetonide (more systemic absorption)

\* For maintenance apply 1-3 times per week

# Why push for otic topical corticosteroids?

Prevent worsening and potentially reverse chronic ear canal changes

## **Reduces new ear infections**

Limits end stage ear disease in predisposed breeds

Improves patient comfort

Chronic ear inflammation is painful and may lead to resistance to ear treatments over time

# Case Presentation-6 year old Poodle Mix (second opinion)

Hx: Ear infections 2-3 times per year since 1 year of age. Since last summer is 'always' having an ear infection. Treated with compounded packing in prior years, but switched to a gentamicin based drop when that stopped working

PE: Both ear canals are firm, but not hard and the patient cries when ear canals are manipulated. Mild salivary staining of groin, otherwise no skin changes. On monthly isoxazoline prevention.



# Case Presentation- 6 year old Poodle Mix

## Second opinion

### **Topical Plan**

Enrofloxacin based ear drop

OR Polymyxin B based ear drop


AND strong topical steroid q 24 hours

AND ear flush containing Triz EDTA q 24-48 hours

### **Systemic Plan**

Oral corticosteroid 0.5-1mg/kg per day until resolution of inflammatory changes, then taper

Gabapentin 10-20mg/kg q 8-12 hours



Mixed population of predominantly rod shaped bacteria

# Case Presentation- 6 year old Poodle Mix

Recheck 3 weeks later

Hx: All oral and ear medications have been administered, patient was a bit resistant of initial therapy but this improved after 1 week. Mild pu/pd.

PE: Both ear canals are firm, but not hard, slight discomfort on palpation. No skin changes.



Vertical canal is smooth, focal erosions of horizontal canal with irregular mass effect in the horizontal ear canals

# Case Presentation- 6 year old Poodle Mix

## Recheck 3 weeks later

### ***Topical Plan***

Enrofloxacin based ear drop

OR Polymyxin B based ear drop


AND fluocinolone acetonide drops q 72 hrs

AND ear flush containing Triz EDTA q 72 hours

### ***Systemic Plan***

Oral corticosteroid 0.5mg/kg q 48 hours x 14 days, then 0.5mg/kg q 72 hours until recheck

Gabapentin 10-20mg/kg q 8-12 hours until recheck



Low number of cocci with few degenerate neutrophils.

# When do I sedate or anesthetize a patient for an ear exam?

Concern for foreign body of mass in the ear canal

**Sedation** for examination and a light cleaning

Neurologic signs

*Heavy rod shaped bacterial overgrowth and initial cleaning may be helpful*

**Anesthesia** for biopsy, most foreign body removals, thorough ear cleaning.

*I need to know the status of the TM prior to dispensing medication*

# Case Presentation- 6 year old Poodle Mix

Recheck 3 weeks later

## Procedure under anesthesia:

Earigator system used to flush debris (does not budge)

Endoscopic forceps grab at material which comes out in large pieces.



‘Mass’ is a mix of packing material, keratinaceous debris, and infectious material

# Case Presentation- 6 year old Poodle Mix

Second recheck 3 weeks later

## ***Topical Plan***

Fluocinolone acetonide drops q 72 hrs

Ceruminolytic ear flush q 72 hrs



## **Systemic Plan**

Consider oclacitinib vs ilunicitinib vs cyclosporine based on response to topical therapy



Opaque tympanic membrane, mild erythema, and mild ceruminous debris

# Brachycephalic external acoustic meatus and TM invisibility



Narrow external acoustic meatus (EAM) and hidden pars tensa in the French bulldog

Mason MJ, Lewis MA. Structure and scaling of the middle ear in domestic dog breeds. *J Anat.* 2024;245(2):324-338. doi:10.1111/joa.14049

# Case Presentation- 4 year old French Bulldog emergency hospital

Hx: Three ear infections in the past year. Environmental allergies treated with lokivetmab q 30 days and intermittent oclacitinib for flares of itch. Body itching responds well to lokivetmab and oclacitinib.

Presents today with concern of eye infection and vomiting

PE: Both ear canals are firm, but not hard and the patient winces when right ear canal is manipulated. Moderately erythematous and slightly swollen interdigital spaces. On monthly isoxazoline prevention.




Mixed ceruminous to purulent golden to green sticky mucoid discharge in the right ear canal.

# 4 year old French Bulldog- neuro examination

## Peripheral vestibular disease and facial nerve paralysis

- Horizontal to Rotary Nystagmus- (fast phase away from the lesion for peripheral vestibular disease)
- Unilateral ventral strabismus
- The right eye is inflamed- corneal ulcer is present
- Complete lack of blink response on the right side



# Case presentation-4 year old French Bulldog


## ***Plan***

Computed Tomography (CT)  
imaging of the skull

Clean ear canal while patient is  
under anesthesia

## ***Topical therapy***

Enrofloxacin based ear drops




Cocci present with nuclear streaming

# Otitis media without bony lysis

## Medical Management




Erythematous and erosive ear canal walls with moderate stenosis and scant purulent debris



Still CT image showing thickening of the tympanic bulla due to bacterial otitis media

# Tympanokeratoma (Cholesteatoma)



Expansile masses of keratin and cholesterol that form secondary to chronic infection or inflammation

More common in brachycephalic dogs

The masses progressively expand leading to lysis of the bulla and neurologic signs.

Surgery is required

# Key Points

- Ear canal inflammation and disease severity does not have to match with skin inflammation and disease severity
- Glucocorticoids topically along with ear canal cleaning is a good first line for uncomplicated otitis externa
- Oral steroids should be considered early in the treatment of chronic ear disease
- Patience is required when treating chronic otitis externa
- The presence of neurologic signs should prompt a recommendation for advanced imaging especially in a brachycephalic breed

Don't just keep up.  
Stay ahead with  
*The Curated Mind*,  
IDEXX's monthly  
education newsletter.

# THE CURATED MIND

CONTINUING  
EDUCATION BROUGHT  
TO YOU BY IDEXX



LEARN EVOLVE REPEAT

**Scan the code** to stay in the know and receive updates about:

- + Upcoming live events
- + New on-demand education resources
- + The latest *Shake Up Your Workup* podcast
- + Articles from *The Vetiverse*
- + Upcoming in-person education opportunities

