




Screening for vector-borne disease

IDEXX 4Dx® Plus Test clinical reference guide



Every dog, every year

The Companion Animal Parasite Council (CAPC) Guidelines recommend annual comprehensive screening for pathogens transmitted by ticks and mosquitoes.¹ Adding annual cycle of comprehensive testing and year-round prevention to your practice benefits your patients, clients, and practice in 3 important ways:

1. React to changing prevalence

Mosquitoes and ticks are constantly on the move, and annual testing is the most reliable way to determine if new infections are threatening pets in your area. Pets move too, of course; without comprehensive testing, you sacrifice the ability to detect and treat mosquito and tick-borne infections acquired in other locations.

2. Detect and treat coinfection

Comprehensive testing lets you assess a dog's risk of having more than one infection.²

3. Measure the effectiveness of prevention protocols

Only comprehensive testing helps you know if your prevention protocols are working. Even a negative result is valuable; it's an opportunity to celebrate the pet owner's role in successfully preventing these infections and keeping their pet healthy.



Know more with every result

With the IDEXX 4Dx® Plus Test, a positive result can also be an indication of ticks and the pathogens they carry.

When you use the IDEXX 4Dx Plus Test as a screening tool, you may

detect antibodies to these pathogens	carried by these ticks	that may also transmit other pathogens and infections to dogs and people	Geographic tick distribution as of 2021 ³
<i>Ehrlichia ewingii</i>	 Lone star tick <i>Amblyomma americanum</i>	<i>Ehrlichia chaffeensis</i> Tularemia Rocky Mountain spotted fever STARI	
<i>Anaplasma phagocytophilum</i> <i>Borrelia burgdorferi</i> (Lyme disease)	 Black-legged tick (deer tick) <i>Ixodes scapularis</i> <i>Ixodes pacificus</i>	<i>Bartonella</i> spp. <i>Babesia</i> spp.	
<i>Ehrlichia canis</i> <i>Anaplasma platys</i>	 Brown dog tick <i>Rhipicephalus sanguineus</i>	<i>Babesia</i> spp. Rocky Mountain spotted fever	
<i>Ehrlichia canis</i>	 American dog tick <i>Dermacentor variabilis</i>	Rocky Mountain spotted fever Tularemia	

Lyme disease

Transmitted by the black-legged tick (deer tick), Lyme disease is caused by the bacterium *Borrelia burgdorferi*. Clinical signs may not appear until several months after infection. Lyme disease has been found throughout North America with cases that have mild to severe disease.

Did you know?

- Dogs testing positive for antibodies to the C₆ peptide had 43% increased risk of having chronic kidney disease (CKD) compared to seronegative dogs.⁴
- The C₆ peptide used in the IDEXX 4Dx[®] Plus Test and Lyme Quant C₆[®] Antibody Test does not cross-react with the antibody response to commercially available Lyme vaccines.⁵
- Dogs with seroreactivity to both *B. burgdorferi* and *Anaplasma phagocytophilum* may have two times the risk of developing clinical illness than singularly infected dogs.²

Borrelia burgdorferi

Primary vectors

Ixodes scapularis or *Ixodes pacificus*
Black-legged tick (deer tick)

Pathology

- Localizes in tissues of infected dogs
- Synovitis (may be subclinical)
- Lyme nephritis

Clinical presentation

Chronic infection with clinical signs that may present acutely:

- Fever, anorexia
- Polyarthritis, lameness
- Rapidly progressive renal failure
- Neurologic syndromes

Laboratory abnormalities

- Elevated C₆ antibody level ≥ 30 U/mL
- May have proteinuria
- May have IDEXX SDMA[®] Test result $> 14 \mu\text{g/dL}$

CKD monitoring

- Chemistry panel with SDMA
 - Recommended to evaluate for the development of protein-losing kidney disease


- Urinalysis with Reflex UPC
 - Recommended to evaluate for proteinuria
- CBC with blood film evaluation
 - Recommended as part of a minimum database



Monitoring considerations

1. If C₆ antibody level drops $\geq 50\%$ from initial level, treatment was successful.
2. If C₆ antibody level drops $< 50\%$ from initial titer, differentials include:
 - a. Noncompliance with treatment. Consider retreating.
 - b. Reinfection. Reevaluate tick control; consider retreating.
 - c. Chronic infection.

*Serology is typically used to diagnose Lyme disease. *B. burgdorferi* localizes to the tissues and is therefore rarely detectable in the blood by PCR.⁶




Heartworm disease

Dirofilaria immitis, the causative agent of heartworm disease, is transmitted when mosquitoes infected with *D. immitis* larvae feed on (or bite) a healthy dog. Heartworm disease has subtle or mild clinical signs in the early stages, making preventive measures so much more important—especially as advanced infection may result in death.

Did you know?

- Despite availability of monthly preventives, prevalence rates of canine heartworm have remained consistent nationwide.⁷
- The American Heartworm Society (AHS) and the Companion Animal Parasite Council (CAPC) recommend testing all dogs for both antigen and microfilariae annually.^{7,8}
- For more information and current recommendations on treating canine heartworm disease, go to heartwormsociety.org or capcvet.org.

<i>Dirofilaria immitis</i>
Primary vector
Mosquito
Pathology
Infective larvae (L3) mature to adult worms in the heart and pulmonary arteries
Clinical presentation
Asymptomatic at first, later developing: • Mild, persistent cough • Lethargy • Exercise intolerance • Reduced appetite • Weight loss
Laboratory abnormalities that may be seen
• Eosinophilia • Azotemia • Increased liver enzymes • Proteinuria



*Administer treatment according to American Heartworm Society protocol.


Canine anaplasmosis

Canine granulocytic anaplasmosis is caused by the bacterium *Anaplasma phagocytophilum* (transmitted by the black-legged tick [deer tick]). *Anaplasma platys* (transmitted by the brown dog tick) is the cause of infectious cyclic thrombocytopenia.

Did you know?

- Many mammalian species, including humans, are susceptible to *A. phagocytophilum* infection.
- Dogs coinfectd with *Anaplasma* and other bacterial pathogens may have more complex disease presentations and respond more slowly to therapy.
- *A. platys* infects canine platelets and is frequently seen as a coinfection with *Ehrlichia canis*.

<i>Anaplasma phagocytophilum</i>		<i>Anaplasma platys</i>	
Primary vectors			
<i>Ixodes scapularis</i> <i>Ixodes pacificus</i> Black-legged tick (deer tick)		<i>Rhipicephalus sanguineus</i> (brown dog tick)	
Pathology			
Infects neutrophils		Infects platelets	
Clinical presentation			
Can present acutely: • Fever • Anorexia • Lethargy • Polyarthritis, lameness • Neurologic signs		Usually minimal clinical signs, but some dogs may have: • Fever • Uveitis • Petechiae and ecchymoses • Epistaxis	
Laboratory abnormalities			
• Thrombocytopenia • Anemia • Lymphopenia • Increased liver enzymes		Other findings may be seen: • Decreased albumin • Increased globulin • Increased ALP and ALT • Proteinuria • Decreased Urine SG • Increased UPC	
Note			
Previous infection may not prevent reinfection and persistent infections are possible. ^{9,10}			



Canine ehrlichiosis

Canine ehrlichiosis is caused by the bacteria *Ehrlichia canis* (transmitted by the brown dog tick) and *Ehrlichia ewingii* (transmitted by the lone star tick). Canine *Ehrlichia* infections may progress to the subclinical phase or may become chronic infections.


Did you know?

- Dogs coinfectd with *E. canis* and *A. platys* were found to have more severe anemia and thrombocytopenia than dogs with either single infection.¹¹
- In a study of healthy dogs with antibodies to *E. canis*, 39% were thrombocytopenic.¹²
- Chronic *E. canis* infections, if left untreated, can lead to bone marrow dysfunction or kidney disease.
- Dogs with *Ehrlichia* antibodies in *E. canis* endemic areas had a 112% increased risk of developing chronic kidney disease (CKD).¹³

CKD monitoring

- Chemistry panel with SDMA
 - Recommended to evaluate for secondary kidney disease.
- Urinalysis with UPC
 - Recommended to evaluate for proteinuria
- CBC with blood film
 - Recommended as part of a minimum database

<i>Ehrlichia canis</i>		<i>Ehrlichia ewingii</i>	
Primary vector			
<i>Rhipicephalus sanguineus</i> (Brown dog tick)		<i>Amblyomma americanum</i> (Lone star tick)	
Pathology			
Infects monocytes		Infects granulocytes	
Clinical presentation			
• Fever, anorexia, lethargy • Bleeding disorders • Polyarthritits, lameness • Lymphadenomegaly • Neurologic signs		• Fever, anorexia, lethargy • Polyarthritits, lameness • Neurologic signs	
Laboratory abnormalities			
• Anemia • Thrombocytopenia • Hyperglobulinemia • Proteinuria			
Other clinical findings may include:			
• Decreased albumin • Increased globulin • Mild increased ALT and ALP • Increased SDMA • Creatinine • Decreased urine specific gravity, proteinuria • Increased urine protein:creatinine (UPC) ratio.			
Note			
Previous infection may not prevent reinfection, and persistent infections are possible. ^{12,14}			





Serology and PCR for sick patients

For dogs presenting with clinical signs consistent with a vector-borne disease, using serology and PCR together improves your ability to make an accurate diagnosis.


Benefits and limitations of each diagnostic method

	Serology	Polymerase chain reaction (PCR)
Measures	Antibody response of host	Nucleic acid (DNA) from pathogen
Benefits	Useful for screening as well as diagnosis of infection	Specifically identifies pathogens indicating active infection
Limitations	Clinical signs may precede a measurable antibody response	A negative PCR result does not necessarily rule out infection

When to use the IDEXX vector-borne disease RealPCR™ panels

- Sick patients with clinical signs and/or laboratory abnormalities consistent with a vector-borne illness
- Patients with subclinical infections based on history, physical examination, serology, and clinical laboratory findings

Dogs with ehrlichiosis and anaplasmosis may present with clinical signs at different times after infection. Which sick dog are you dealing with?



“No single test is sufficient for diagnosing an infectious disease in a sick patient.”

Edward Breitschwerdt, DVM, DACVIM*
Professor, Internal Medicine
College of Veterinary Medicine,
North Carolina State University

*Dr. Breitschwerdt has a business relationship with IDEXX pursuant to which he receives compensation from IDEXX from time to time. The views expressed in this guide are solely those of Dr. Breitschwerdt.

Depend on the most accurate and comprehensive screen

Available in-clinic or from IDEXX Reference Laboratories



SNAP 4Dx Plus Test


Reference-laboratory quality in the palm of your hand, for superior diagnostic accuracy at the point of care.



Lab 4Dx Plus Test

Available from IDEXX Reference Laboratories as a stand-alone test or as part of a more comprehensive parasite screen with the Fecal Dx Antigen Panel with Lab 4Dx Plus Test-Canine.

SNAP® technology uses a proprietary three-step process to deliver dependable sensitivity and specificity.



References

1. General guidelines: Parasite testing and protection guided by veterinarians [dog]. Companion Animal Parasite Council website. www.capvet.org/guidelines/general-guidelines. Updated July 29, 2020. Accessed November 17, 2021.
2. Beall MJ, Chandrashekar R, Eberts MD, et al. Serological and molecular prevalence of *Borrelia burgdorferi*, *Anaplasma phagocytophilum*, and *Ehrlichia* species in dogs from Minnesota. *Vector-Borne Zoonotic Dis*. 2008;8(4):455–464. doi:10.1089/vbz.2007.0236
3. Regions where ticks live [maps]. Centers for Disease Control and Prevention website. www.cdc.gov/ticks/geographic_distribution.html. November 17, 2021.
4. Drake C, Coyne M, McCrann DJ, Buch J, Mack R. Risk of development of chronic kidney disease after exposure to *Borrelia burgdorferi* and *Anaplasma* spp. *Top Companion Anim Med*. 2020;42:100491. doi:10.1016/j.tcam.2020.100491
5. O'Connor TP, Esty KJ, Hanscom JL, Shields P, Philipp MT. Dogs vaccinated with common Lyme disease vaccines do not respond to IR₆, the conserved immunodominant region of the VlsE surface protein of *Borrelia burgdorferi*. *Clin Diagn Lab Immunol*. 2004;11(3):458–462. doi:10.1128/CDLI.11.3.458-462.2004
6. Straubinger RK. PCR-based quantification of *Borrelia burgdorferi* organisms in canine tissues over a 500-day postinfection period. *J Clin Microbiol*. 2000;38(6):2191–2199. doi:10.1128/JCM.38.6.2191-2199.2000
7. CAPC prevalence maps: heartworm [dog]. Companion Animal Parasite Council website. www.capvet.org/maps/#/2021/all-year/heartworm-canine/dog/united-states. Accessed November 17, 2021.
8. American Heartworm Society. *Current canine guidelines for the prevention, diagnosis, and management of heartworm infection in dogs*. 2020. Accessed November 17, 2021. https://d3f8b8ckhnm2.cloudfront.net/images/pdf/AHS_Canine_Guidelines_11_13_20.pdf?1605565616
9. Egenvall A, Lilliehöök I, Björnsdóttir A, et al. Detection of granulocytic *Ehrlichia* species DNA by PCR in persistently infected dogs. *Vet Rec*. 2000;146(7):186–190. doi:10.1136/vr.146.7.186
10. Breitschwerdt EB, Hegarty BC, Qurollo BA, et al. Intravascular persistence of *Anaplasma platys*, *Ehrlichia chaffeensis*, and *Ehrlichia ewingii* DNA in the blood of a dog and two family members. *Parasit Vectors*. 2014;7:298. doi:10.1186/1756-3305-7-298
11. Gaunt S, Beall M, Stillman B, et al. Experimental infection and co-infection of dogs with *Anaplasma platys* and *Ehrlichia canis*: hematologic, serologic and molecular findings. *Parasit Vectors*. 2010;3(1):33. doi:10.1186/1756-3305-3-33
12. Hegarty BC, de Paiva Diniz PP, Bradley JM, Lorentzen L, Breitschwerdt E. Clinical relevance of annual screening using a commercial enzyme-linked immunosorbent assay (SNAP 3Dx) for canine ehrlichiosis. *J Am Anim Hosp Assoc*. 2009;45(3):118–124. doi:10.5326/0450118
13. Burton W, Drake C, Ogeer J, et al. Association between exposure to *Ehrlichia* spp. and risk of developing chronic kidney disease in dogs. *J Am Anim Hosp Assoc*. 2020;56(3):159–164. doi:10.5326/JAAHA-MS-7012
14. Starkey LA, Barrett AW, Beall MJ, et al. Persistent *Ehrlichia ewingii* infection in dogs after natural tick infestation. *J Vet Intern Med*. 2015;29(2):552–555. doi:10.1111/jvim.12567
15. Snellgrove AN, Krapinaya I, Ford SL, et al. Vector competence of *Rhipicephalus sanguineus* sensu stricto for *Anaplasma platys*. *Ticks Tick Borne Dis*. 2020;11(6):101517. doi:10.1016/j.ttbdis.2020.101517



Strengthen the bonds.®



IDEXX Laboratories, Inc.

One IDEXX Drive
Westbrook, Maine 04092
United States

American dog tick (*Dermacentor variabilis*) photographer: Susan E. Ellis, USDA-APHIS-PPQ. Black-legged tick (*Ixodes scapularis*), lone star tick (*Amblyomma americanum*), and brown dog tick (*Rhipicephalus sanguineus*) photographer: James L. Occi.

© 2022 IDEXX Laboratories, Inc. All rights reserved. • 09-69074-13

All ®/TM marks are owned by IDEXX Laboratories, Inc. or its affiliates in the United States and/or other countries.

The IDEXX Privacy Policy is available at [idexx.com](https://www.idexx.com).